

International Journal of Modern Physics A
© World Scientific Publishing Company

J/Ψ DISSOCIATION AND ADLER'S THEOREM

ALEX BOURQUE

*Department of Physics, McGill University
3600 University Street, Montreal, QC, Canada H3A 2T8
bourquea@physics.mcgill.ca*

CHARLES GALE

*Department of Physics, McGill University
3600 University Street, Montreal, QC, Canada H3A 2T8
bourquea@physics.mcgill.ca*

KEVIN L. HAGLIN

*Department of Physics & Astronomy, Saint Cloud State University
720 Fourth Avenue South, St. Cloud, MN 56301, USA
klhaglin@stcloudstate.edu*

Effective Lagrangian models of charmonium have recently been used to estimate dissociation cross sections with light hadrons. Detailed study of the symmetry properties reveals possible shortcomings relative to chiral symmetry. We therefore propose a new Lagrangian and point out distinguishing features amongst the different approaches. Using the newly proposed Lagrangian, which exhibits $SU_L(4) \times SU_R(4)$ symmetry and complies with Adler's theorem, we find dissociation cross sections with pions that are reduced in an energy dependent way, with respect to cases where the theorem is not fulfilled.

Keywords: charmonium; QGP; Adler's theorem.

PACS numbers: 11.25.Hf, 123.1K

1. Introduction

The theoretical study of matter under extreme conditions enjoys a wide range of application, from the physics of the early Universe, to that of relativistic nuclear collisions. The latter offer the tantalizing possibility of recreating in the laboratory the conditions that prevailed roughly a microsecond after the Big Bang. The theory of the strong interaction, Quantum Chromodynamics (QCD), predicts a phase transition from normal hadronic matter to a plasma of quarks and gluons¹. To find a signature of this new state of matter represents a task which has generated tremendous activity both in theory and in experiment. Of the many signals put forward as probes of the quark-gluon plasma, the suppression of the J/ψ yield enjoys a popular status².

2 *A. Bourque, C. Gale, and K. L. Haglin*

Indeed, since charmonium is predominantly produced in the early stage of the nuclear collisions through hard processes, it acts as a probe for the subsequent stages. The original idea was that the presence of a quark-gluon plasma (QGP) will screen the long-range confining force between $c\bar{c}$, leading to the decoherence of the pair². This suppression mechanism was later augmented by the possibility of charmonium dissociation by hard gluons in a deconfined medium³. But, in order to identify the true nature of a possibly new phase, it appears necessary to quantify the J/ψ -light hadron interaction. This requirement on the J/ψ cross-sections with light mesons is not a trivial one to satisfy, as there are no direct experimental measurements. One has to rely on theoretical calculations based, for example, on QCD sum-rules⁴, on quark-potential models^{5,6}, on extended Nambu and Jona-Lasinio models^{7,8}, or on effective SU(4) mesonic Lagrangians^{9,10,11,12,13}. These lead typically to cross-sections from a few tenths of a millibarn to a few millibarns near threshold⁴.

Specifically, for the effective mesonic Lagrangians found in Refs 11, 12, 13, once form factors have been folded-in to account for short-range interactions, the dissociation cross-section by pions reaches a few millibarns. But concerns have been raised about using such models: (i) the SU(4) symmetry used to describe the pseudoscalar and vector meson interactions is questionable as it is broken, (ii) the form factors accounting for the finite size of the mesons do not proceed from the formalism as in other models^{4,6}, and (iii) the $J/\psi + \pi \rightarrow D^* + \bar{D}$ process does not vanish in the soft-pion limit for non-degenerate vector meson masses as expected from Adler's theorem¹⁴. In this contribution we will outline the consequence of implementing the soft-pion theorem (i.e., Adler's theorem) on the dissociation cross-section of $J/\psi + \pi \rightarrow D^* + \bar{D}$ by constructing a Lagrangian invariant under the global $SU_L(4) \times SU_R(4)$ symmetry group.

2. Adler's theorem

A spontaneously broken symmetry not only implies the existence of Goldstone bosons, but also constrains their low-energy behaviour. Here we will consider the transition amplitude for emitting one Goldstone boson (the proof can easily be extended to more than one). Following Weinberg¹⁵, we first note that the current operator can create a Goldstone boson

$$\langle 0 | J^\mu(x) | B \rangle = i \frac{F q^\mu}{(2\pi)^3 2E} e^{-iq \cdot x}, \quad (1)$$

where F is the decay constant. We expect the matrix element $\langle \beta | J^\mu(x) | \alpha \rangle$ to then have a pole term. Indeed, in general, we can write

$$\langle \alpha | J^\mu(0) | \beta \rangle = N_{\beta\alpha}^\mu + i \frac{F q^\mu}{q^2} M_{\beta\alpha}^B, \quad (2)$$

where $M_{\beta\alpha}^B$ is the desired transition amplitude for emitting one Goldstone boson, and $N_{\beta\alpha}^\mu$ is assumed to be the pole-free contribution. Applying the current conser-

vation constraint, we then find that

$$M_{\beta\alpha}^B = \frac{i}{F} q_\mu N_{\beta\alpha}^\mu \quad (3)$$

and we see that as $q \rightarrow 0$ the RHS vanishes. Historically, this property was first studied by Adler¹⁶, and rests on the assumption that $N_{\beta\alpha}^\mu$ has no singularity as $q \rightarrow 0$. This is not valid in general, since a pole may arise through the insertion of the current operator on an external line¹⁷. $M_{\beta\alpha}^B$ can then have a non-vanishing contribution in the soft limit due to emission of a soft Goldstone boson off an external line. But for chiral-symmetric and normal-parity mesonic Lagrangians such an insertion is impossible and we expect the theorem to hold. We will now explicitly check that decoupling occurs for the process $\pi^+ + \rho^0 \rightarrow \pi^+ + \rho^0$ in two chiral-invariant Lagrangians.

2.1. Chiral model with π , ρ , and a_1 mesons

We first build the effective chiral Lagrangian as in Ref. 18 yielding

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} \text{Tr} [\partial_\mu \phi \partial^\mu \phi] - \frac{1}{4} \text{Tr} [F_{\mu\nu}^V F_V^{\mu\nu} + F_{\mu\nu}^A F_A^{\mu\nu}] \\ & + \frac{1}{2} m_V^2 \text{Tr} [V_\mu^2] + \frac{1}{2} m_A^2 \text{Tr} [A_\mu^2] \\ & + \mathcal{L}_{V\phi\phi} + \mathcal{L}_{AV\phi} + \mathcal{L}_{VV\phi\phi} + \cdots, \end{aligned} \quad (4)$$

where ϕ is the pseudo-scalar field, F_V and F_A are the field strength tensors for the vector and axial vector mesons respectively. The model has three parameters, namely the pion decay constant, F_π , the vector mass, m_V , and the universal coupling constant, g . The axial vector meson mass is related to the vector mass in this model and thus is not an input parameter.

It is then seen that the process under consideration, namely $\pi^+(p_1) + \rho^0(q_1) \rightarrow \pi^+(p_2) + \rho^0(q_2)$, involves 5 tree-level diagrams for the given particle content: π - and a_1 -mediated exchanges both in s and t channels, and through a 4-point interaction (Fig 1).

Extracting the relevant interaction terms results in the five off-shell amplitudes


$$i\mathcal{M}_{\mu\nu}^1 = i\Gamma_\mu^{1\dagger} \frac{i}{s - m_\pi^2} i\Gamma_\nu^1, \quad (5)$$

$$i\mathcal{M}_{\mu\nu}^2 = i\Gamma_\mu^{1\dagger} \frac{i}{t - m_\pi^2} i\Gamma_\nu^1, \quad (6)$$

$$i\mathcal{M}_{\mu\nu}^3 = i\Gamma_{\mu\alpha}^{2\dagger} \frac{-i[g^{\alpha\beta} - (p_1 + q_1)^\alpha (p_1 + q_1)^\beta / m_A^2]}{s - m_A^2} i\Gamma_{\nu\beta}^2, \quad (7)$$

$$i\mathcal{M}_{\mu\nu}^4 = i\Gamma_{\mu\alpha}^{2\dagger} \frac{-i[g^{\alpha\beta} - (q_2 - p_1)^\alpha (q_2 - p_1)^\beta / m_A^2]}{t - m_A^2} i\Gamma_{\nu\beta}^2, \quad (8)$$

$$i\mathcal{M}_{\mu\nu}^5 = i\Gamma_{\mu\nu}^3, \quad (9)$$

4 *A. Bourque, C. Gale, and K. L. Haglin*Fig. 1. $\rho^0 + \pi^+ \rightarrow \rho^0 + \pi^+$.

where the vertex functions are

$$\Gamma_\mu^1 = -\frac{g}{\sqrt{2}} \left[k_\mu + p_\mu - \frac{(1-Z^2)}{m_V^2} (q \cdot k p_\mu - q \cdot p k_\mu) \right], \quad (10)$$

$$\Gamma_{\mu\nu}^2 = -\frac{ig}{\sqrt{2}} \frac{(1-Z^2)^{\frac{1}{2}}}{m_V} \{ (m_A^2 + q^2 - k^2) g_{\mu\nu} - q_\mu q_\nu + k_\mu k_\nu \}, \quad (11)$$

$$\Gamma_{\mu\nu}^3 = 2 \left[\frac{g^2}{2} \frac{(1-Z^2)}{2m_V^2} \{ -2p_1 \cdot p_2 g_{\mu\nu} + p_{1\mu} p_{2\nu} + p_{2\mu} p_{1\nu} \} + \frac{g^2}{2Z^2} g_{\mu\nu} \right], \quad (12)$$

for $\pi^+(p) + \rho^0(q) \rightarrow \pi^+(k)$, $\pi^+(p) + \rho^0(q) \rightarrow a_1^+(k)$, and the four-point interaction, respectively, and the shorthand $Z^2 = \left(1 - \frac{g^2 \tilde{F}_\pi^2}{4m_0^2} \right)$.

Having now the full amplitude for the given process, we wish to see if the pseudoscalar decoupling theorem holds. The presence of the contact term in the 4-point interaction leads us to expect cancellations to occur as we let one of the pions' 4-momentum go to zero. Stated differently, since the ρ -meson was introduced in a chirally symmetric way by adding its chiral partner (i.e. a_1), the transition amplitude relies on help from the a_1 in what amounts to a delicate cancellation allowing the pion to decouple. To show this, we contract the amplitudes with the appropriate polarization vectors, and then let $p_2 \rightarrow 0$ (a similar proof can be shown to hold for $p_1 \rightarrow 0$). First it is seen that the amplitudes involving the π -exchange go to zero because of transversality (i.e $\epsilon(q) \cdot q$). For a_1 -exchange in the s -channel we find (the

same result is true for the t -channel)

$$\begin{aligned}
i\mathcal{M}_3 &= \epsilon^{*\mu}(q_2) \left[i\Gamma_{\mu\alpha}^{2\dagger}(p_2 \rightarrow 0) \frac{-i \left[g^{\alpha\beta} - \frac{q_2^\alpha q_2^\beta}{m_A^2} \right]}{m_V^2 - m_A^2} i\Gamma_{\nu\beta}^2 \right] \epsilon^\nu(q_1) \\
&= -\frac{ig^2}{2} \frac{1}{m_V^2} \epsilon^{*\mu}(q_2) \left[m_A^2 g_{\mu\nu} - q_{1\mu} q_{1\nu} + \frac{q_1 \cdot q_2}{m_A^2} q_{2\mu} q_{1\nu} + \frac{m_V^2}{m_A^2} q_{2\mu} q_{2\nu} \right] \epsilon^\nu(q_1) \\
&= -\frac{ig^2}{2Z^2} \epsilon^*(q_2) \cdot \epsilon(q_1).
\end{aligned} \tag{13}$$

The last line comes about again due to the orthogonality condition. Finally, the 4-point interaction reads

$$i\mathcal{M}_5 = \frac{ig^2}{Z^2} \epsilon^*(q_2) \cdot \epsilon(q_1), \tag{14}$$

and thus the *full amplitude* is shown to vanish as expected. Note the cancellation between the 4-point interaction and the a_1 channels. In summary, even though the pions are not coupled through gradient coupling for all interaction terms, the net amplitude still vanishes due to intricate cancellations amongst all the channels.

2.2. Chiral model with π and ρ mesons

In the previous section the axial mesons were introduced as the chiral partners of the ρ fields resulting in a linear realization of the symmetry¹⁹. In the present case, since the desired chiral Lagrangian will involve only pseudoscalar and vector mesons, both will then have to transform non-linearly under the symmetry. This is done by gauging-away the axial mesons²⁰ and yields

$$\begin{aligned}
\mathcal{L} &= -\frac{1}{4} \text{Tr} [F_{\mu\nu}(\rho) F^{\mu\nu}(\rho)] + \frac{1}{2} m_V^2 \text{Tr} [\rho_\mu^2] \\
&+ \frac{F_\pi^2}{4} (1+k) \text{Tr} [\partial_\mu \zeta \partial^\mu \zeta] + i \frac{F_\pi^2 g_{V\phi\phi}}{2} \text{Tr} [\rho^\mu (\partial_\mu \zeta \zeta^\dagger + \partial_\mu \zeta^\dagger \zeta)] \\
&+ \frac{F_\pi^2}{4} (1-k) \text{Tr} [\zeta^\dagger \partial^\mu \zeta^\dagger \zeta \partial_\mu \zeta],
\end{aligned} \tag{15}$$

where the input parameters are now m_V^2 , $g_{V\phi\phi}$, F_π , and k , and the field strength tensor is $F_{\mu\nu}(\rho) = \partial_\mu \rho_\nu - \partial_\nu \rho_\mu - ig[\rho_\mu, \rho_\nu]$.

In this model there are only two diagrams, namely s - and t -channels of pion exchange. The relevant interaction Lagrangian is

$$\mathcal{L}_{\rho\pi\pi} = -i \frac{g_{\rho\pi\pi}}{2} \text{Tr} [\rho^\mu [\phi, \partial_\mu \phi]], \tag{16}$$

and the extracted vertex for the s - and t - channels is

$$\Gamma_\mu'^1 = -\frac{g_{\rho\pi\pi}}{\sqrt{2}} (p_\mu + k_\mu), \tag{17}$$

$$\tag{18}$$

6 *A. Bourque, C. Gale, and K. L. Haglin*

with $k = p + q$ and $k = p - q$, respectively. The two amplitudes are then

$$i\mathcal{M}_{\mu\nu}^1 = i\Gamma_{\mu}^{\prime 1\dagger} \frac{i}{s - m_{\pi}^2} i\Gamma_{\nu}^{\prime 1}, \quad (19)$$

$$i\mathcal{M}_{\mu\nu}^2 = i\Gamma_{\mu}^{\prime 1\dagger} \frac{i}{t - m_{\pi}^2} i\Gamma_{\nu}^{\prime 1}. \quad (20)$$

We immediately see that the soft pion theorem holds separately for each amplitude when the proper polarization vectors are contracted with the amplitudes.

3. Application to the J/Ψ dissociation

We first start by noting that all the effective Lagrangian models found in the literature ^{11,12,13} are invariant under $SU_V(4)$ (in the degenerate mass limit), but not under the extended chiral symmetry, i.e., $SU_L(4) \times SU_R(4)$, and therefore we expect the pions will not decouple in soft-momentum limit. Specifically, here we will compare the $J/\psi + \pi \rightarrow D^* + \bar{D}$ cross-section as calculated within the $SU_V(4)$ -only models found in Refs 11, 12, 13 and with the Lagrangian of Eq. (15) extended to the $SU_L(4) \times SU_R(4)$ symmetry group. For the $SU_V(4)$ models, three diagrams need to be considered (Fig. 2), while for the extended chiral model proposed here, the contact interaction is absent. It can be explicitly shown that the pseudoscalars decouple in the soft momentum limit for that latter case ²¹.




Fig. 2. $J/\psi + \pi \rightarrow D^*(\bar{D}^*) + \bar{D}(D)$.

The differential isospin-averaged cross-section is given by

$$\frac{d\sigma}{dt} = \frac{1}{128\pi s p_1^2} \mathcal{M}^{\mu\nu} \mathcal{M}^{\alpha\beta} \left[g_{\mu\alpha} - \frac{q_{2\mu} q_{2\alpha}}{m_{D^*}^2} \right] \left[g_{\nu\beta} - \frac{q_{1\nu} q_{1\beta}}{m_{J/\psi}^2} \right], \quad (21)$$

where the appropriate model-dependent squared amplitude is used, an isospin factor

of two has been included and the centre of mass momentum is

$$p_1^2 = \frac{1}{4s} \lambda(s, m_\pi^2, m_{J/\psi}^2) \quad (22)$$

and the triangle function is $\lambda(x, y, z) = x^2 - 2x(y + z) + (y - z)^2$. Integrating over the kinematical range defined by

$$t_\pm = m_\pi^2 + m_{D^*}^2 - \frac{1}{2s}(s + m_\pi^2 - m_{J/\psi}^2)(s + m_{D^*}^2 - m_D^2) \pm \frac{1}{2s} \lambda^{1/2}(s, m_\pi^2, m_{J/\psi}^2) \lambda^{1/2}(s, m_D^2, m_{D^*}^2) \quad (23)$$

gives the total cross-section. Fixing the relevant parameters²¹ and carrying this to completion for the two models yields Fig. 3. We see an energy-dependent reduction in the cross sections across the relevant domain and to quote a specific number we note that at $\sqrt{s} = 5$ GeV the cross-section is reduced by about 40% going from the $SU_V(4)$ model to the $SU_L(4) \times SU_R(4)$ model.




Fig. 3. Isospin-averaged cross-section for $J/\psi + \pi \rightarrow (D^* + \bar{D}) + (\bar{D}^* + D)$.

4. Conclusion

Modeling low-energy hadron physics is particularly challenging when there is limited experimental input available for constraints. This is the current situation in the charm sector as the only measurement relevant for fixing coupling constants in the model is the decay width for $D^* \rightarrow D\pi$ ²². We have therefore invoked symmetries and general theorems. In particular, we have checked for full $SU_L(4) \times SU_R(4)$ symmetry and the appropriate limit to test for compliance with Adler's theorem. We found that none of the published models can do this, and we therefore proposed a new effective Lagrangian—the first one which does encode complete four-flavour chiral symmetry and Adler's theorem. Our interest here has been solely to quantify the effect of these. A complete calculation including form factors and a longer list of reactions is a topic for a separate study.

Since Adler's theorem is relevant at low-energy, the near-threshold cross sections are expected to be affected the most. We found the cross section for $J/\psi + \pi \rightarrow (D^* + \bar{D}) + h.c.$ to be reduced as compared to a choice of Lagrangian which does not encode the full flavour chiral symmetry and does not obey Adler's theorem. The reduction is energy dependent, but seems to be a few tens of percents from threshold to $\sqrt{s} = 5$ GeV. In a full calculation the size of this reduction might not persist when one takes into account not only form factors, but also abnormal parity interactions and symmetry breaking effects (e.g. pseudoscalar masses and non-degenerate vector mass spectrum).

Abnormal parity interactions may play an important role near threshold. Indeed, it was shown that Adler's theorem breaks down if a soft Goldstone boson can be inserted on an external line. This is expected to happen for abnormal parity Lagrangians where a $VV\phi$ vertex exists. The abnormal parity contribution to the $J/\psi + \pi$ amplitude will then not vanish in the soft limit. The problem in including these interactions lies again in the lack of experimental data to fix the coupling strengths.

Symmetry breaking effects are also expected to be important since the underlying $SU_L(4) \times SU_R(4)$ is broken.

It will also be important in the future to take this formalism to completion by implementing covariant hadronic form factors computed within the same effective Lagrangian or perhaps other approaches. Ultimately, the outlook for this line of study is to estimate the dissociation cross sections with all the light hadrons, with finite size effects incorporated, and then to input the results into a dynamical model for heavy ion reactions to finally address the question of J/ψ survivability in the hadronic phase (primarily mesonic matter). For then, one would have a more complete understanding of the J/ψ yield and therefore know what it implies about QGP formation.

Acknowledgments

A.B. thanks S. Turbide for helpful discussions, and A.B. and C.G. thank E. S. Swanson for a useful visit. This work was supported in part by the Natural Sciences and Engineering Research Council of Canada, in part by the Fonds Nature et Technologies of Quebec, and in part by the National Science Foundation under grant number PHY-0098760.

References

1. F. Karsch and E. Laermann, *Quark-Gluon Plasma 3*, eds Rudolph C. Hwa and Xin-Nian Wang (World Scientific, Singapore, 2004).
2. T. Matsui and H. Satz, *Phys. Lett.* **B178**, 416 (1986).
3. D. Kharzeev and H. Satz, *Phys. Lett.* **B334**, 155 (1994).
4. D. Kharzeev, H. Satz, A. Syamtomov, and G. Zinovjev, *Phys. Lett.* **B389**, 595 (1996); F. O. Duraes, H. C. Kim, S. H. Lee, F. S. Navarra, and M. Nielsen, *Phys. Rev.* **C68**, 035208 (2003).
5. K. Martins, D. Blaschke, and E. Quack, *Phys. Rev.* **51**, 2723 (1995).
6. C.-Y. Wong, E.S. Swanson, and T. Barnes, *Phys. Rev.* **C65**, 014903 (2001).
7. M.A. Ivanov, J. G. Körner, and P. Santorelli, *Phys. Rev.* **D70**, 014005 (2004).
8. A. Deandrea, G. Nardulli, and A. D. Polosa, *Phys. Rev.* **D68**, 034002 (2003).
9. S. G. Matinyan and B. Müller, *Phys. Rev.* **C51**, 2723 (1995).
10. K. L. Haglin, *Phys. Rev.* **C61**, 031902 (2000).
11. K. L. Haglin and C. Gale, *Phys. Rev.* **C63**, 065201 (2001).
12. Z. Lin, C.M. Ko, and B. Zhang, *Phys. Rev.* **C61**, 024904 (2000).
13. Y. Oh, T. Song, and S. Houn Lee, *Phys. Rev.* **C63**, 034901 (2001).
14. F. S. Navarra, M. Nielson, and M. R. Robilotta, *Phys. Rev.* **C64**, 021901 (2001).
15. S. Weinberg, *The Quantum Theory of Fields*, Vol. 2 (Cambridge University Press, 1995).
16. S. Adler, *Phys. Rev.* **137**, B1022 (1965).
17. J.J. Sakurai, *Currents and Mesons* (University of Chicago Press, Chicago, 1969).
18. U. Meissner, *Phys. Rept.* **161**, 215 (1988).
19. S. Gasiorowicz and D. A. Geffen, *Rev. Mod. Phys.* **41**, 531 (1969).
20. J. Schechter, *Phys. Rev.* **D34**, 868 (1986).
21. A. Bourque, C. Gale, H. L. Haglin, *Phys. Rev.* **C70**, 055203 (2004).
22. A. Anastassov *et al.*, *Phys. Rev.* **D65**, 032003 (2002).